

Critical exponents for random knots

Alexander Yu. Grosberg,^{1,2,3,4}

¹*Department of Physics and Center for Materials Science and Engineering,
Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

⁴*On leave from: Institute of Biochemical Physics, Russian Academy of Sciences, Moscow 117977, Russia*

The size of a zero thickness (no excluded volume) polymer ring is shown to scale with chain length N in the same way as the size of the excluded volume (self-avoiding) linear polymer, as N^ν , where $\nu \approx 0.588$. The consequences of that fact are examined, including sizes of trivial and non-trivial knots.

Knots are encountered in virtually every branch of physics, as evidenced by the series of books on “Knots and Everything.” Knots keep entertaining physicists and mathematicians for one and a half century, since W.Thomson [1]. In a more specific context, knotted polymers and DNA remain in the center of attention for over three decades [2–4]. Nevertheless, many simple basic questions remain unanswered. For instance, the most fundamental physical property of any macromolecule - its average size $R(N)$, scaling with chain length N , and depending on the solvent conditions - is not understood for knotted polymers. In particular, the value of gyration radius determines diffusion, light scattering, gel penetration and several other properties of macromolecules. It is known in details for a variety of linear polymers under different solvent conditions. The mere closing ends and making a ring-shaped chain instead of a linear one reduces its size only marginally, by a factor of $\sqrt{2}$ for Gaussian chain and certainly preserving the power law even for excluded volume one. The really strong dramatic effect, as we show below in this paper, is due to the quenched restricted set of conformations - those which are topologically equivalent. Indeed, topological type must be quenched for all real closed ring-shaped polymers whose segments are unable to pass through each other. The only exception is DNA in the presence of topo-II enzyme [5], and the very fact that Nature has invented such marvelous tool indicates how serious is the topological problem for long polymers. Thus, the formulation of the problem is very simple: given the polymer ring of a quenched topology, we want to estimate its average size. So far, there have been only very few works on this problem [6,7], and overall the problem remains poorly understood. The purpose of this note is to propose some heuristic arguments addressing this problem.

1. In this paper, we consider the simplest ring polymer with no excluded volume [8], consisting of some N freely-jointed straight segments, length ℓ each. Assume that every spatial shape, or conformation, of this polymer is just as likely as any other. In mathematical language, our “polymer” is simply a closed broken line embedded in 3D. With probability 1, this object does not have self-intersections, and, therefore, represents a knot. It may be a trivial knot, if it can be continuously and without self-

crossings transformed into a plane polygon topologically equivalent to a circle, or it may be one of the non-trivial knots. It has been proven as early as in 1988 [10] that the probability of a trivial knot configuration decays exponentially with the chain length, N :

$$p_{\text{tk}}(N) \simeq \exp(-N/N_0) , \quad (1)$$

where “tk” subscript stays for “trivial knot” and a constant N_0 represents the characteristic length. This mathematical prediction has been beautifully confirmed in computer simulation [11], yielding $N_0 \simeq 340 \pm 4$.

The restriction to broken-line polymer of straight segments is not necessary. Alternatively, one can think, for instance, of a smooth worm-like ring object of the length $N\ell$ with effective segment ℓ [12]. One can also make straight segments with Gaussian distributed random lengths [13]. The exponential result (1) holds for these models as well as for all similar ones, except perhaps with different values of N_0 . However, one cannot use continuous model (the Wiener trajectory), since it corresponds to the limit $N \rightarrow \infty$, $\ell \rightarrow 0$, $N\ell = \text{const}$, where essentially all conformations are heavily knotted, or trivial knot probability is exactly zero.

While rigorous proof of the result (1) is not simple, it can be readily understood on a hand-waving level. Indeed, consider our polymer as a sequence of N/g blobs, with g segments in each blob. The probability of trivial topology for each blob is $p_{\text{tk}}(g)$, and formula (1) is equivalent to saying that $p_{\text{tk}}(N) = [p_{\text{tk}}(g)]^{N/g}$, which means that topological constraints between blobs are negligible compared to that between segments inside each blob. The temptation is then to conclude that *vast majority of knots occur on the small length scales*. However appealing, this jargon statement is confusing: indeed, while blob may be much smaller than the entire polymer, $g \ll N$, the argument above only holds if it is large compared to N_0 ($g \gg N_0$), which itself is numerically large.

Exponentially small probability of a trivial knot (1) immediately explains why topological constraints *can* significantly alter the polymer size. Indeed, $\ell N^{1/2}$ is the size averaged over *all* conformations. However, the size of a trivial knot is an average over an exponentially small *subset* of conformations. When relatively compact complex knots are removed, the remaining average increases. In

order to bring this idea to a quantitative level, a couple of further preliminary arguments will be handy.

2. Consider now two ring polymers, with N segments each. They form a link, trivial or non-trivial. If rings overlap, i.e. the distance between their centers of mass is smaller than their sizes, the probability that the link is trivial vanishes, $p_{\text{tl}}(N) \rightarrow 0$, when $N \rightarrow \infty$. While the exact asymptotic expressions for $p_{\text{tl}}(N)$ are not known, it should be qualitatively similar to the trivial knot probability (1), and we assume that the corresponding characteristic chain length N_1 is of the order of N_0 .

Speaking about the trivial link probability, one has to make the distinction between a link made by two rings each of which may be an arbitrary knot, and a link made by two trivially knotted rings. Since the former probability is higher, and goes to 0 for large N , the latter, which we use below, goes to zero as well.

3. Since trivial link topology is highly unlikely for overlapping rings, the untangled ring polymers effectively present excluded volume for each other, even if there is no bare excluded volume for monomers. Simple scaling argument suggests that the excluded volume can be estimated as

$$v_{\text{excl}}(N) \simeq R^3(N) [1 - p_{\text{tl}}(N)] , \quad (2)$$

where $R(N)$ is the (properly averaged) size of one ring. The concept of “topological excluded volume” was introduced in one of the first works on computer simulation of knots and links [15]. It can be easily understood in terms of an exactly solved model [16,17] in which one polymer is a ring and the other is a straight line. It has recently been rigorously proven [18] that the excluded volume scales as $R^3(N)$ in $N \rightarrow \infty$ limit for two rings if their Gaussian linking number remains zero, even though this condition does not guarantee the untangled topology. As regards eq. (2), it is actually almost trivially correct. Indeed, $1 - p_{\text{tl}}(N)$ is the probability for two phantom rings to be tangled, i.e., to adopt a conformation prohibited for real (non-phantom) untangled rings [19].

4. We are now prepared to directly attack the question of the size, $R_{\text{tk}}(N)$, of a trivial knot without excluded volume. Let us color two different pieces of our N -segment ring, both containing some $g \ll N$ monomers (segments). Since neither of the pieces has open ends, they present uncrossable objects for one another. Therefore, the arguments above apply and suggest that these two pieces exclude for each other some volume which is of the order of $v_{\text{excl}}(g)$. While this identification of the pieces of a ring chain with separate small rings may be doubtful for small g , we expect it to become increasingly accurate with increasing g , especially when g becomes greater than N_0 . In that case, $p_{\text{tl}}(g)$ is negligible, and we end up with $v_{\text{excl}}(g) \simeq R^3(g)$ for $g > N_0$, which immediately implies that the chain of g -blobs belongs to the universality class of self-avoiding walks, and thus

$$R_{\text{tk}}(N) \simeq R(g) (N/g)^\nu , \quad \text{where } \nu \approx 0.588 \approx 3/5 . \quad (3)$$

We expect this to hold at $g > N_0$, while at smaller scales the effect of topological constraints is only marginal. Thus, we assume $R(g) \simeq \ell g^{1/2}$ for g up to about N_0 and apply the above result (3) choosing $g \sim N_0$:

$$R_{\text{tk}}(N) \simeq \begin{cases} \ell N^{1/2} & \text{if } N < N_0 \\ \ell N_0^{-\nu+1/2} N^\nu & \text{if } N \gg N_0 \end{cases} . \quad (4)$$

Thus, chain size in the case of trivial knot topology is controlled by the standard excluded volume exponent close to $3/5$. This confirms the conjecture made by J. des Cloizeaux as early as in 1981 [6]. We see also that this “swollen” regime comes only for really long chains, because N_0 is numerically large [11]. Furthermore, while in terms of parameters the cross-over to the “swollen” asymptotics is controlled by N_0 , there are grounds to expect that this asymptotics actually develops at N around a few times N_0 (which we expressed with the $\gg$ sign in formula (4)). On the other hand, although topological constraints lead to swelling only when N is really large, the prefactor in the second line of formula (4) is of order unity: $N_0^{-\nu+1/2} \approx 340^{-0.088} \approx 0.6$.

It is an interesting challenge to produce a renormalization group analysis for the trivial knot polymer. One of the difficulties is that regular ϵ -expansion techniques are hardly applicable, since topological constraints do not exist for linear polymers in dimensions other than 3; as a matter of fact, these constraints exist in the space of dimension d for the objects of dimension $d-2$. The manifestation of that difficulty is seen in the fact that trivial link probability enters the expression (2) for the scale-dependent excluded volume. On the other hand, such RG calculation, once achieved, can shed a long awaited light on why N_0 is as large numerically as it is.

5. Consider now some non-trivial knot, still without any excluded volume. This can be addressed using the so-called “tube inflation” or “ideal knot” approach. This idea has been proposed in several contexts by different authors, in the aspect addressing knot entropy it was suggested in [20]. Recently, the approach received a significant attention [21]. In particular, the present author’s contribution in this book addressed the size of an arbitrary knot. However, the result (4) was not established at the time; instead, the size of an arbitrary knot was estimated under the assumption that size of a trivial knot scales as N^μ with an unknown exponent μ . We can now directly use the results of the author’s work in [21], simply substituting $\mu = \nu$ there. To make the present work self-contained, we repeat the argument briefly.

When polymer is a knotted ring, fluctuations of each segment are restricted by the neighboring segments. Following the standard trick of the reptation theory [22], we replace these constraints by an effective tube confining the entire polymer. The difficult part of the problem is that, unlike the usual case of a melt [22], the tube shape for the swollen single chain fluctuates very strongly, and we have to find a reasonable approximation for it. To this end, we argue that there are two possibilities to achieve

maximal entropy, corresponding to “uniform” and “phase segregated” structures. Let us examine both.

We call the structure “uniform” when the freedom of transverse fluctuations is about the same everywhere along the polymer. That corresponds to finding the maximal diameter, D , for which the tube of the length L can be knotted into the given knot without self-penetration. The shape of this “maximally inflated” tube, or the shape of its central axis line, is called “ideal representation of a knot.” It is characteristic of the knots topology, and so is this tube “axis ratio” $p = L/D$.

Thus, we replace the real topological constraints with the confinement within the tube whose aspect ratio, p , is that characteristic for the given knot topology. Tube is closed, and polymer makes exactly one turn within the tube. We assume further that the polymer is not knotted inside the tube [23]. These two conditions together guarantee that the polymer has exactly the desirable topology. We can now determine overall size of the polymer in space, R , using the following Flory-style argument: When R increases, polymer loses entropy because it gets stretched along the tube axis; when R decreases, polymer loses entropy because it gets squeezed perpendicular to the tube axis. Thus, the equilibrium size R can be conjectured to balance these two entropic factors. To implement this idea, we choose D and L such that $LD^2 \sim R^3$ and, since $L/D = p$, that means $L \simeq Rp^{2/3}$, $D \simeq Rp^{-1/3}$. We further recall that the unknotted polymer behaves just as an excluded volume one on large scales. That allows us to employ the standard scaling arguments [24] to estimate the relevant entropy. Specifically, transverse compression and longitudinal stretching in the tube are associated with, respectively, concentration blobs and Pincus tension blobs:

$$S \simeq - \left(\frac{L}{N^\nu} \right)^{1/(1-\nu)} - \left(\frac{N^{3\nu}}{LD^2} \right)^{1/(3\nu-1)}. \quad (5)$$

Provided the above expressions for L and D , this entropy has a maximum with respect to R at $R \sim \ell N^\nu p^{-\nu+1/3}$, and we identify it as the average size of the knot. This result applies as long as each of the aforementioned blobs remains larger than N_0 monomers. Equivalently, equilibrium tube diameter D should be greater than ℓN_0^ν , yielding $N > pN_0$. For smaller blobs, Gaussian statistics should be valid, which means physically that we can consider a phantom [19] polymer confined within the tube. This case has been already considered in [20]. Collecting everything together, we obtain

$$R_k(N) \simeq \begin{cases} \ell N^{1/2} p_k^{-1/6} & \text{if } N < p_k N_0 \\ \ell N_0^{-\nu+1/2} N^\nu p_k^{-\nu+1/3} & \text{if } N > p_k N_0 \end{cases}, \quad (6)$$

where we have included the subscript in p_k to emphasize that this parameter is taken for the given knot k .

Thus, in terms of N -dependence in the limit of very large N , every knot acts as an effective excluded volume, making the exponent equal to $\nu \approx 0.588$. However, more

complex knots, with larger p , are significantly less expanded than trivial or simple knots. Furthermore, for complex knots the onset of swollen behavior is pushed to the range of really long chains. These results are consistent with numerical observation [11,25] suggesting that probability of any particular knot, and not only trivial one, decays exponentially at large N (1), with the characteristic length N_0 independent of the knot type. Qualitatively, one can say that increasing N for the polymer with any given topology eventually leads to the situation when polymer is dominated by a very long strings in which knot restriction is not felt at all, and thus it should become in this sense equivalent to the trivial knot.

Another way to look at the eq. (6) is from the point of view of p dependence. For the very long chain, with $N \gg N_0$, topological “memory” acts as an excluded volume for relatively simple knots, with $p < N/N_0$, for which R obeys the last line of eq. (6). With increasing knot complexity, the system crosses over to the other regime, described by the upper line in eq. (6). Note that at the cross-over the chain size is about $R \sim \ell N^{1/3} N_0^{1/6}$. In terms of N -dependence, this is already a collapsed chain.

Note also that the maximal value of entropy (5) is about $S \sim -p$: the knot creates about one constraint per polymer length which is about tube diameter.

6. As we mentioned, there is an alternative way to maximize entropy, which we call “phase segregated.” To explain what we mean by that, consider first a prime knot. It may be entropically favorable to tighten this knot as much as possible, losing virtually all the freedom for segments involved in the tight knot, but gaining maximal entropy for the rest of the chain which then fluctuates as a free unknotted loop. For the composite knot, this picture must be slightly modified: we still imagine a long free unknotted loop with several little knots - prime components of the original composite knot - independently tightened in different places on the loop.

To estimate roughly the entropy of such knot segregated state, we can use again formula (5), applying it twice (or several times for a composite knot): for the tightened part, where p is roughly equal to that of the original knot, and for the unknotted loop part, where p is of order one. The chain lengths for these two are about p and $N - p$, respectively. Since equilibrium entropy is linear in p , this simplest estimate yields that phase segregated state is about as favorable entropically as the uniform one. A more sophisticated approach may be needed to decide which of them is more stable. It is also likely that the answer is sensitive to the details of local chain geometry (eg, freely-jointed vs worm-like segments, and the like).

7. Although the qualitative statement that $p_{\text{tl}}(N)$ is practically zero for long rings is sufficient for the main stream of arguments in this paper, to estimate $p_{\text{tl}}(N)$ is itself an interesting problem. The simple estimate is obtained in the following way. Consider two strongly overlapping rings. Since we are counting probabilities over all

possible configurations of rings, regardless of their knottedness, the dominant statistics is Gaussian, and there should be about $\sqrt{N}$ contacts between the rings. As long as N is not very large, each contact can be thought of as capable to provide up to one (positive or negative) unit of linking. Therefore, there are about $\sqrt{N}$ different linking states, with the probability of order $1/\sqrt{N}$ for each of them, including the untangled one. When N gets really very large, the situation becomes more complicated, because each of the $\sqrt{N}$ “contacts” is itself a large crowd of segments, which can realize a variety of linking arrangements. Since entanglements in different contact regions do not commute to each other in this regime [14], there appears exponential diversity of different types of linkings. Adjusting prefactors for continuity, we arrive at

$$p_{\text{tl}}(N) \sim \begin{cases} \sqrt{N_1/N} & \text{if } N < N_1 \\ \exp \left[1 - \sqrt{N/N_1} \right] & \text{if } N > N_1 \end{cases}, \quad (7)$$

where subscript “tl” stays for the trivial link, N_1 is the characteristic length. The first line of this formula is in a very good agreement with the simulation data of the work [25] where links of the lengths up to 500 were examined. This is consistent with our conjecture that N_1 is of the order of N_0 . It would be interesting to simulate longer chains, as we expect the N -dependence to cross-over to exponential in the range of several hundred segments.

It is also interesting to present the blob-style argument to explain the lower line of the eq. (7). Consider each ring as consisting of N/g blobs, with g segments in each blob. About $\sqrt{N/g}$ of the blobs from one ring are expected to overlap with the blobs of the other ring. Now the bold assumption comes (supported by an admittedly vague argument): in order for two rings not to be linked, all of the blobs must be untangled. Since the “dangerous” blobs are only the overlapping ones, we obtain $p_{\text{tl}}(N) = [p_{\text{tl}}(g)]^{\sqrt{N/g}}$, which yields the familiar answer (7).

When this work has already been written, the paper [7] appeared. It reports simulation data for the ring-shaped polymer which preserves the topology of a trivial knot and does not have any excluded volume. The average size of this polymer has been measured for a series of lengths $N = 2^n$, ranging from 32 to 2048 monomers. The data are consistent with our results. Specifically, the sizes of N -mers were found to follow Gaussian statistics, $N^{0.5}$, for small N ($n = 5, 6, 7$), then power was slowly increasing with N , and finally the ratio of the sizes of 2048- and 1024-mers was close to $2^{0.58}$. This seems in perfect agreement with the result eq. (4).

The author has a pleasure to acknowledge useful discussions with B.Duplantier, I.Eruxhimovich, S.Nechaev, D.Nelson, T.Witten.

-
- [1] Kelvin, Lord *Transactions of the Royal Society of Edinburgh*, v. **25**, p. 217 - 260, 1868.
 [2] M.Delbrück *Mathematical Problems in the Biological Sciences, Proceedings of Symposium in Applied Mathematics*

- (American Mathematical Society, Providence RI), v. **14**, p. 55-68, 1962.
 [3] H.L.Frisch, E.Wasserman *J. Am. Chem. Soc.*, **83**, p. 3789, 1961.
 [4] ref about DNA knots,
 [5] Sikorav J.-L., B. Duplantier, G. Jannink & Y. Timsit. *J. Mol. Biol.* v. **284**, p. 1279-1287, 1998; *J. Mol. Biol.* v. **284**, p. 1289-1299, 1998.
 [6] J. des Cloizeaux *J. Physique Lettres*, v. **42**, p. L433, 1981.
 [7] J.M.Deutsch *Phys. Rev. E*, v. **59**, n. 3, p. R2539, 1999.
 [8] Excluded volume can be neglected under the condition $z = (d/\ell)\sqrt{N} \ll 1$ [9], where d is the segment diameter. Under this condition volume exclusion is completely insignificant, while the very fact that segments cannot pass through one another is preserved, and so is the topology
 [9] A.Grosberg, A.Khokhlov *Statistical Physics of Macromolecules*, AIP press, NY, 1994 (page 91).
 [10] D.W.Summers, S.G.Whittington *Journal of Physics A: Math.& Gen.*, v. **21**, p. 1689, 1988; N.Pippenger *Disc. Appl. Math.*, v. **25**, 273, 1989.
 [11] T.Deguchi, K.Tsurusaki *Phys. Rev. E*, v. **55**, n. 5, p. 6245-6248, 1997.
 [12] Worm-like chain adopts smoothly curved conformations $\mathbf{r}(s)$ with the weight proportional to $\exp [-(\ell/2) \int \dot{\mathbf{r}}^2 ds]$, where s is arc length along the chain, $\ddot{\mathbf{r}} = d^2\mathbf{r}/ds^2$, integration runs along the entire polymer length $N\ell$, and ℓ is called the effective Kuhn segment.
 [13] K.Koniaris, M.Muthukumar *Phys. Rev. Lett.*, v. **66**, p. 2211-2214, 1991.
 [14] Nechaev, S. *Statistics of knots and entangled random walks*, (World Scientific: Singapore, 1996)
 [15] A.V.Vologodskii, A.V.Lukashin, M.D.Frank-Kamenetskii *Sov. Phys. JETP*, v. **40**, p. 932-936, 1975.
 [16] S.F.Edwards *Proceedings of the Physical Society*, v. **91**, p. 513, 1967.
 [17] S.Prager, H.Frisch *Journal of Chemical Physics*, v. **46**, p. 1475, 1967.
 [18] B.Duplantier, private communication.
 [19] We adopt here the terminology in which *phantom* stays for the chain whose segments can freely pass through one another, irrespectively of excluded volume.
 [20] A.Yu.Grosberg, A.Feigel, Y.Rabin *Physical Review E*, v. **54**, n. 6, p. 6618-6622, 1996.
 [21] V.Katrich, A.Stasiak, L.Kauffman, Editors *Ideal Knots*, World Scientific, 1999.
 [22] M.Doi, S.F.Edwards *The Theory of Polymer Dynamics*, Oxford University Press, Oxford 1986.
 [23] The concept of an unknotted polymer confined within knotted tube is pretty subtle. It can be explained in the following way. First, let us quench our polymer inside the tube with respect to the tube walls. Then, let us pretend that the tube is phantom and bring it to the trivial knot non-overlapping state. If our polymer is in a trivial knot state after this procedure, we say that it was unknotted within the tube.
 [24] P.G. de Gennes *Scaling Concepts in Polymer Physics*, Cornell Univ.Press, Ithaca, NY, 1979.
 [25] T.Deguchi, K.Tsurusaki *Lectures at Knots96*, edited by S.Suzuki, World Scientific, 1997, p. 95.